

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Archives of Cytology and Histopathology Research

Journal homepage: <https://www.achr.co.in/>

Original Research Article

Correlation of radiological parameters with cytological finding in the diagnosis of thyroid swelling

Amar Kumar¹, Manish Kumar^{1,*}, Swati Sucharita Giri¹, Dharitri Rabha¹, Richa¹¹Dept. of Pathology, Patna Medical College and Hospital, Patna, Bihar, India

ARTICLE INFO

Article history:

Received 10-11-2021

Accepted 03-12-2021

Available online 04-03-2022

Keywords:

Thyroid malignancy

Ultrasound parameters

Cytology

ABSTRACT

Introduction: Thyroid gland diseases are the most common endocrine disease seen in clinical practice after Diabetes mellitus. Ultrasound parameters along with correlation with cytology finding helps in differentiate between benign and malignant thyroid nodule.

Aim: The purpose of this study was to study the sonographic features of various benign and malignant thyroid nodules, and to correlate the sonographic findings with Fine Needle Aspiration Cytology (FNAC) and so as to evaluate the accuracy of ultrasonography in diagnosing benign and malignant nodules.

Materials and Methods: The study was conducted in the Pathology department in collaboration with the Radiology Department on 209 patients who were send for ultrasound for thyroid swelling. These patients were further send for fine needle aspiration to the pathology department.

Result: The majority of cases which turned malignant on cytology report in this study have association with ultrasound features like predominant solid lesion, hyperechoic lesion, irregular margin, AT ratio greater than 1, absent peripheral halo, microcalcification along with increased internal vascularity and solitary nodules.

Conclusion: Ultrasonography of thyroid nodules along with fine needle aspiration cytology serves as a best screening test to detect malignancy in outpatient department. The use of different parameters in ultrasound helps in categorizing the lesion and their management. In this study we have association with ultrasound features like predominant solid lesion, hyperechoic lesion, irregular margin, AT ratio less than 1, absent peripheral halo, microcalcification along with increased internal vascularity and solitary nodules with malignancy on cytology.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Thyroid gland disease are the most common endocrine disease seen in clinical practice after Diabetes mellitus.¹ The prevalence of thyroid related disease is 3-8 % of the general population contributing around 42 million people in india are suffering from thyroid disease presently in India as per various studies.^{2,3} Ultrasonography (USG) gives good knowledge of its internal anatomy and its relation with adjacent organs and details of different

pathological features which occurs in benign and malignant lesion of thyroid without using ionizing radiation or iodine-containing contrast medium.^{4,5} It tells about internal composition (solid or cystic), presence of nodularity, echogenecity of mass, invasion in nearby structures, antero-transverse diameter, assessment of blood flow pattern in and around lesion, calcification and presence of peripheral halo to differentiate between benign and malignant thyroid nodule.^{6,7} Fine-needle aspiration cytology (FNAC) is the first investigation of choice in thyroid swellings. Fnac is simple and quick to perform in the outpatient department has excellent patient compliance, and can be repeated

* Corresponding author.

E-mail address: amarkumarsah2009@gmail.com (M. Kumar).

in case of doubt.⁸ FNAC is the best single test for discriminating malignant thyroid nodules due to its high sensitivity and specificity, Innumerable research have been performed for use in thyroid ultrasonography (USG) parameters to differentiate benign from malignant thyroid nodules.⁹⁻¹³

2. Aim of the study

The purpose of this study was to study the sonographic features of various benign and malignant thyroid nodules, and to correlate the sonographic findings with Fine Needle Aspiration Cytology (FNAC) and so as to evaluate the accuracy of ultrasonography in diagnosing benign and malignant nodules.

3. Materials and Methods

The study was conducted in the Pathology department in collaboration with the Radiological Department on 209 patients who were send for ultrasound for thyroid swelling. These patients were further send for fine needle aspiration to the pathology department.

3.1. Inclusion criteria

1. Patients selected for this study were undiagnosed cases with palpable thyroid swelling.
2. Patients selected for this study were undiagnosed cases with clinically non palpable thyroid swelling but detected on ultrasound.

3.2. Exclusion criteria

1. Patients excluded were from previously diagnosed cases of thyroid disease.
2. Patients excluded were follow up for treatment of thyroid disease.

The investigations were performed on cases using a high frequency probe ultrasound machine. The 8 parameters used were

1. Internal Composition (Solid, Predominantly solid, Cystic, Predominantly cystic and Spongiform)
2. Echogenecity (isoechoic, hyperechoic, hypoechoic, heterogeneous)
3. Margins (Well defined or Poorly defined)
4. Antero-posterior and Transverse Ratio (AT Ratio > 1 or < 1)
5. Peripheral halo (Present or Absent)
6. Calcification (Macro-calcification or Micro-calcification)
7. Internal Vascularity (increased or decreased or peripheral)
8. Nodules (Absent or single or multiple)

Fine needle aspiration was done using a 21 gauge spinal needle with suction using 10 ml syringe under all aseptic conditions. The cytology slides were air dried and wet fixed in Absolute Ethanol and air dried were stained with Liesman – Giemsa stain and wet fixed slides were stained with Papanicolaou stain. These slides were examined under microscope and categorize into Benign and Malignant on cytology and were subclassified also.

4. Result

In this study it was seen that out of 209 patients 83.73 % of patients are female and 16.26% were male. The mean Age (Years) was 36.67 ± 15.17 .

In this study it was seen most patients were in the age group of 40-49 yrs(24%) followed by 20-29 yrs (23%) in the female group while in the male patient most patients are in age group of 20-29 yrs(33%) followed by 40-49 yrs and 50-59 yrs (18%).

Among 209 thyroid cases, 15 cases were reported malignant in Fnac report. Fisher's exact test was used to explore the association between cytological diagnosis and ultrasound parameters.

Among the Ultrasound features regarding composition, predominantly Solid (20.0%) had the largest proportion of Malignant report on cytology while predominantly cystic, cystic, spongiform had the largest proportion of Benign on cytology. Regarding echogenecity, hyperechoic (11.1%) and hypoechoic (7.0%) had strongest association with malignant on cytology report while hypoechoic (93%) on ultrasonography were associated with benign on cytology report.

In respect to margin on ultrasound, irregular margin (20.6%) on USG had the largest proportion of malignant on cytology report while defined Margin (98.6%) of the patients were turned benign on cytology report. Among the ultrasound features regarding AT Ratio, AT Ratio: > 1 had the largest proportion of association with Malignant on cytology and Antero-Transverse Ratio: < 1 had the largest proportion of association with Benign on cytology.

In respect to peripheral halo, absent peripheral halo (8.1%) were associated more with malignancy on cytology and peripheral halo present (97.2%) were more associated with benign on cytology. In view of calcification observed in this study, Microcalcification was strongly associated with malignancy on fnac report and Macrocalcification were strongly associated with benign on cytology report.

Among the internal vascularity features observed in ultrasound, increased internal vascularity (22.8%) were strongly associated more with malignancy on cytology and peripheral vascularity (100%) was associated more with benign on cytology. Among types of nodules, solitary nodules (8.0%) are associated more with malignancy on cytology while multiple nodules (92.5%) are associated more with benign on cytology report.

The majority of cases which turned malignant on cytology report in this study have association with ultrasound features like predominant solid lesion, hyperechoic lesion and irregular margin, AT ratio more than 1, absent peripheral halo, microcalcification along with increased internal vascularity and solitary nodules.




Fig. 1: Showing benign thyroid follicular cells in monolayered sheets in Colloid goiter with background of thin colloid. (MGG Stain 100X).




Fig. 2: Thyroid follicular cells showing open fine chromatin and intranuclear inclusion (MGG Stain 400X).

5. Discussion

The present study was focused on correlation of ultrasound features with cytological diagnosis has shown similarity of parameters with the other studies. Most of the cases of thyroid nodules were in age group of 20-29 in males and 40-49 in females.

Thus in this study we have seen that a predominantly solid nodule, hyperechogenicity, irregular margin, AT Ratio > 1 and microcalcification and increased internal vascularity




Fig. 3: USG showing a well margined solitary anechoic cystic colloid nodule




Fig. 4: USG showing a well defined homogeneous solitary thyroid nodule




Fig. 5: USG showing Solitary inhomogeneous, hypoechoic mass lesion, irregular in outline and showing multiple punctate echogenicity representing microcalcifications (psammoma bodies)

Table 1: Gender distribution

	Number	Percentage
Female	175	83.73%
Male	34	16.26%
Total number of Cases	209	100%

Table 2: Age and Gender wise Distribution of Cases.

Age group	Female		Male		Grand Total
<20	23	13%	2	6%	12%
20-29	41	23%	11	33%	25%
30-39	36	21%	3	9%	19%
40-49	42	24%	6	18%	23%
50-59	17	10%	6	18%	11%
60-69	16	9%	4	12%	10%
70-79	00	0%	1	3%	0%
Total	175	100%	34	100%	100%

Table 3: Distribution of the Cases in Terms of Different Ultrasound features.

USG Features	Benign	Malignant	Total
Composition			
Predominantly Solid	24	6	30
Solid	82	08	90
Predominantly Cystic	38	0	38
Cystic	49	0	49
Spongiform	02	0	02
Echogenecity			
Isoechoic	14	01	15
Hyperechoic	25	02	27
Hypoechoic	132	10	142
Heterogenous	24	01	25
Margins			
Defiened	145	01	146
Poorly defiened	40	13	63
At ratio			
AT Ratio <1	166	09	175
AT Ratio >1	29	05	34
Peripheral HALO			
Present	35	01	36
Absent	160	13	173
Calcification			
Macrocalcification	41	01	42
Microcalcification	08	04	12
Absent	146	09	155
Internal vascularity			
Increased	44	13	57
Decreased	140	01	141
Pheripheral	11	00	11
Nodules			
Absent	14	00	14
Solitary	82	06	88
Multiple	101	06	107

Table 4: Distribution of the cases in terms of cytological features

Cytological Features	Frequency	Percentage
Colloid Goiter	123	58.9%
Hyperplastic Nodule	22	10.5%
Lymphocytic Thyroiditis	15	7.2%
Hashimoto's Thyroiditis	12	5.7%
Papillary Thyroid Carcinoma	9	4.3%
Thyroglossal Cyst	5	2.4%
Adenomatoid Nodule	4	1.9%
Benign Cystic Lesion	3	1.4%
Follicular Neoplasm	3	1.4%
Aberrant Thyroid	2	1.0%
Adenomatoid Goiter	2	1.0%
De Quervian's Thyroiditis	2	1.0%
Thyroid Adenoma	2	1.0%
Anaplastic Carcinoma	1	0.5%
Chronic Lymphocytic Thyroiditis	1	0.5%
Follicular Thyroid Carcinoma	1	0.5%
Giant Cell Thyroiditis	1	0.5%
Medullary Thyroid Carcinoma	1	0.5%

Table 5: Sensitivity and Specificity and Positive and Negative Predictive value along with Accuracy of different USG Parameters

Parameters	Sensitivity	Specificity	PPV	NPV	Diagnostic Accuracy
Predominant Solid	100.0%	45.9%	12.5%	100.0%	49.8%
Hypochoic/Hyperechoic	86.7%	19.6%	7.7%	95.0%	24.4%
Irregular Margin	86.7	74.2%	20.6%	98.6%	75.1%
Antero-Transverse Ratio > 1	40.0%	85.6%	17.6%	94.9%	82.3%
Absent Peripheral Halo	93.3%	18.0%	8.1%	97.2%	23.4%
Microcalcification	26.7%	95.9%	33.3%	94.4%	90.9%
Internal vascularity	11.0%	94.3%	0.0%	92.4%	87.6%
Solitary Nodules	53.3%	49.0%	7.5%	93.1%	49.3%

have more diagnostic accuracy to detect malignancy on cytology.

In our study among USG parameters like internal composition we have noticed predominant solid lesion has more association with malignancy, having sensitivity of 100% and specificity of 45% with accuracy of 49.8%. Sharma et al¹⁴ had reported a solid lesion has high sensitivity (100%) in predicting malignancy but accuracy is low (49.2%). Frates et al¹⁵ also reported that solid composition has highest sensitivity (of 69.0% to 75.4%) in predicting malignancy; however the chance of being malignant of solid nodule predictive value is low (15.6% - 27%) Kwak et al¹⁶ also reported solid echotexture has more association with malignancy.

Regarding echogenicity in our study we have hyperechoic followed by hypoechoic both are associated with malignancy having sensitivity of 86 % and specificity of 19.6% with accuracy of 24.4%. Moon et al¹⁷ reported that a hypoechoic nodule had a sensitivity of 87.2%, specificity of 58.5% and an accuracy of 70.7% in predicting malignancy while Sharma et al¹⁴ had reported a sensitivity of 85.7%, specificity of 67.5% and an accuracy of 69.5%

in predicting malignancy. Few studies however showed result inconsistent with the literature, hypoechogenicity and the presence of hypoechoic rim did not affect the risk of malignancy.

In respect to margin in our study, Irregular margin were associated more with malignancy having sensitivity of 86.7% and specificity of 74.7% with accuracy of 75.2%. Sharma et al¹⁴ showed poorly defined margins have sensitivity of 78.5%, specificity of 82.2% and a diagnostic accuracy of 81.8%. Hoang et al¹⁸ reported sensitivity of ill-defined margins ranges from 53%–89%. Therefore, unless frank invasion beyond the capsule (if more than 50% of its border is not clearly demarcated) is demonstrated, the US appearance of the nodule margins alone is an unreliable basis for determining malignancy.

Malignant nodules often assume a taller-than-wider shape, i.e., antero-posterior diameter > transverse diameter on a Ultrasound. We reported sensitivity of 40 % and specificity of 85.6 % with accuracy of 82.3 % for AT ratio > 1 in detecting malignancy. Cappelli et al¹⁹ showed sensitivity of 99 % and specificity of 57 % in detecting malignancy. Kim et al²⁰ found that a solid thyroid nodule

AT ratio > 1 has 93% specificity for malignancy. Sharma et al¹⁴ showed specificity of 87% and the highest diagnostic accuracy of 87.5% for diagnosing a malignant nodule.

An incomplete or complete absence of peripheral halo is often associated with a malignant nodule. Our study demonstrated that the absent peripheral halo sign had a sensitivity of 93.3% and specificity of 18.0 % with accuracy of 23.4% indicating that it is only a low predictor malignancy. Sharma et al¹⁴ showed sensitivity of 64.2% and an accuracy of 54.3% in their study and while Rago et al²¹ showed absent peripheral halo had sensitivity of 66.6% and specificity of 77%.

In other studies microcalcification served as best predictor of malignancy. Pallaniappan et al²² reported that microcalcification had 100% specificity for malignancy, which is similar to our study. Hoang et al²³ stated that microcalcification are one of the most specific features of thyroid malignancy, with a specificity of 85.8%–95% and a positive predictive value of 41.8%–94.2% We got sensitivity of 26.7% and specificity of 95.9% with accuracy of 90.9% which is similar to other studies.

Chan et al²³ reported that study had some intrinsic blood flow is seen in malignancy, and they concluded that completely avascular nodule is very unlikely to be malignant. Sharma et al¹⁴ stated increased internal vascularity seen in malignant nodules with a sensitivity of 85.7% and an accuracy of 66.6%. Our study also showed 11.0% sensitivity and specificity of 94.3% with accuracy of 87.3%.

In view of nodularity, our study showed sensitivity of 53.3% and specificity of 49.0% with accuracy of 49.3% with solid nodules. Ugurlu et al²⁴ study having a single nodule or two nodules increased the chance of malignancy which showed consistency with our study. But Taneri et al²⁵ reported that multiple nodules in thyroid glands were associated with malignancy.

6. Conclusion

Ultrasonography of thyroid nodules along with fine needle aspiration cytology serves as a best screening test to detect malignancy in outpatient department. The use of different parameters in ultrasound helps in categorizing the lesion and their management. In this study we have association with ultrasound features like predominant solid lesion, hyperechoic lesion and irregular margin, AT ratio more than 1, absent peripheral halo, microcalcification along with increased internal vascularity and solitary nodules with malignancy on cytology.

7. Conflict of Interest

The authors declare that there is no conflict of interest.

8. Source of Funding

None.

References

1. Rojeski MT, Gharib H. Nodular thyroid disease. Evaluation and management. *N Engl J Med.* 1985;313(7):428–36. doi:10.1056/NEJM198508153130707.
2. Wiest PW, Hartshorne MF, Inskip PD, Crooks LA. Thyroid palpation versus high-resolution thyroid ultrasonography in the detection of nodules. *J Ultrasound Med.* 1998;17(8):487–96.
3. Morstensen JD, Woolner LB, Bennett WA. Gross and microscopic findings in clinically normal thyroid glands. *J Clin Endocrinol Metab.* 1955;15(10):1270–80. doi:10.1210/jcem-15-10-1270.
4. Butch RJ, Simeone JF, Mueller PR. Thyroid and parathyroid ultrasonography. *Radiol Clin North Am.* 1985;23(1):57–71.
5. Leopold GR. Ultrasonography of superficially located structures. *Radiol Clin North Am.* 1980;18(1):161–73.
6. Clark KJ, Cronan JJ, Scola FH. Color Doppler sonography: Anatomic and physiologic assessment of the thyroid. *J Clin Ultrasound.* 1995;23(4):215–23.
7. Foley WD, Erickson SJ. Color Doppler flow imaging. 1991;156(1):3–13. doi:10.2214/ajr.156.1.1898567.
8. Baloch ZW, Livolsi VA. Fine-needle aspiration of the thyroid: Today and tomorrow. *Best Pract Res Clin Endocrinol Metab.* 2008;22(6):929–39. doi:10.1016/j.beem.2008.09.011.
9. Grant CS, Hay ID, Gough IR, McCarthy PM, Goellner JR. Long-term follow-up of patients with benign thyroid fine needle aspiration cytologic diagnoses. *Surgery.* 1989;106(6):980–5.
10. Gharib H, Goellner JR. Fine needle aspiration biopsy of the thyroid: an appraisal. *Ann Internal Med.* 1993;118(4):282–9. doi:10.7326/0003-4819-118-4-199302150-00007.
11. Gharib H. Fine-needle aspiration biopsy of thyroid nodules: advantages, limitations, and effect. *Surgery.* 1994;69(1):44–9. doi:10.1016/s0025-6196(12)61611-5.
12. Walfish PG, Hazani E, Strawbridge H, Miskin M, Rosen B. A prospective study of combined ultrasonography and needle aspiration biopsy in the assessment of the hypofunctioning thyroid nodule. *Surgery.* 1977;82(4):474–82.
13. Miller JM, Hamburger JI, Kini S. Diagnosis of thyroid nodules. Use of fine-needle aspiration and needle biopsy. *J Am Med Assoc.* 1979;241(5):481–4.
14. Sharma G. Ultrasonographic Evaluation of Thyroid Nodules with Pathologic Correlation. *Int J Anatomy.* 2017;6(2):53–7.
15. Frates MC, Benson CB, Charboneau JW, Cibas ES, Clark OH, Coleman BG, et al. Management of thyroid nodules detected at US: Society of Radiologists in Ultrasound Consensus Conference Statement. *Radiology.* 2005;237(3):794–800. doi:10.1148/radiol.2373050220.
16. Kwak JY, Han KH, Yoon JH, Moon HJ, Park SH, Jung HK, et al. Thyroid imaging reporting and data system for US features of nodules: a step in establishing better stratification of cancer risk. *Radiology.* 2011;260(3):892–9. doi:10.1148/radiol.11110206.
17. Moon WJ, Jung SL, Lee JH, Na DG, Baek JH, Lee YH, et al. Benign and malignant thyroid nodules: US differentiation - multicenter retrospective study. *Radiology.* 2008;247(3):762–70.
18. Hoang JK, Lee WK, Lee M, Johnson D, Farrell S. US features of thyroid malignancy. Pearls and Pitfalls. *Radiographics.* 2007;27(3):847–65. doi:10.1148/rg.273065038.
19. Cappelli C, Castellano M, Pirola I, Gandossi E, Martino E, Agosti DCB, et al. Thyroid nodule shape suggests malignancy. *Eur J Endocrinol.* 2006;155(1):27–31. doi:10.1530/eje.1.02177.
20. Kim EK, Park CS, Chung W, Oh KK, Kim DI, Lee JT, et al. New sonographic criteria for recommending fine-needle aspiration biopsy of nonpalpable solid nodules of the thyroid. *AJR Am J Roentgenol.* 2002;178(3):687–91.
21. Rago T, Vitti P, Chiovato L, Mazzeo S, Liperi AD, Miccoli P. Role of Conventional Ultrasonography and Color Flow-Doppler Sonography in Predicting Malignancy In ‘Cold’ Thyroid Nodules. *Eur J Endocrinol.* 1998;138(1):41–6.

22. Palaniappan MK, Aiyappan SK, Ranga U. Role of gray scale, color Doppler and spectral Doppler in differentiation between malignant and benign thyroid nodules. *J Clin Diag Res.* 2016;10(8):1–06.
23. Chan BK, Desser TS, McDougall IR, Weigel RJ, Jr RBJ. Common and uncommon sonographic features of papillary thyroid carcinoma. *J Ultrasound Med.* 2003;22(10):1083–90. doi:10.7863/jum.2003.22.10.1083.
24. Ugurlu S, Caglar E, Yesim TE, Tanrikulu E, Can G, Kadioglu P, et al. Evaluation of thyroid nodules in Turkish population. *Intern Med.* 2008;47(4):205–9.
25. Taneri F, Kurukahvecioglu O, Ege B, Yilmaz U, Tekin E, Cifter C, et al. Prospective analysis of 518 cases with thyroidectomy in Turkey. *Endocr Regul.* 2005;39(3):85–90.

Author biography

Amar Kumar, Tutor

Manish Kumar, Associate Professor  <https://orcid.org/0000-0003-1752-8509>

Swati Sucharita Giri, Tutor

Dharitri Rabha, Junior Resident

Richa, Junior Resident

Cite this article: Kumar A, Kumar M, Giri SS, Rabha D, Richa. Correlation of radiological parameters with cytological finding in the diagnosis of thyroid swelling. *IP Arch Cytol Histopathology Res* 2022;7(1):9-15.