

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Archives of Cytology and Histopathology Research

Journal homepage: <https://www.achr.co.in/>

## Case Report

# Kikuchi-Fujimoto disease in a young male post-COVID -19 vaccination

Jackson Bose<sup>1</sup>, Jamunarani Srirangaramasamy<sup>1</sup>, Khowsalya Subrajaa Karupasamy<sup>1</sup>,  
Sekar Manjani<sup>1\*</sup>

<sup>1</sup>Dept. of Pathology, Bhaarith Medical College and Hospital (BIHER), Chennai, Tamil Nadu, India



### ARTICLE INFO

#### Article history:

Received 05-12-2023

Accepted 21-12-2023

Available online 08-01-2024

#### Keywords:

Granulomatous inflammation

Radiological imaging

COVID -19

### ABSTRACT

Kikuchi disease (Kikuchi-Fujimoto disease) is one of the rarest forms of lymphadenopathy. It is also known as necrotizing lymphadenitis. The patients usually present with fever and lymphadenopathy. Here is a case of Kikuchi-Fujimoto disease presented with fever and cervical lymphadenopathy. Radiological imaging revealed matted lymph nodes. Fine needle aspiration of the lymph node revealed granulomatous inflammation. Acid Fast stain was negative and further investigations ruled out tuberculosis. A histopathological biopsy of the lymph node confirmed the diagnosis of Kikuchi-Fujimoto disease. The patient was treated with medications and the patient's condition improved in follow-up. Tuberculosis must be ruled out in such conditions as the symptoms and radiological and FNAC findings mimicking tuberculosis. The diagnosis of Kikuchi-Fujimoto disease is made by histopathology and confirmed by immunohistochemistry.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

Kikuchi-Fujimoto disease is one of the rarest forms of lymphadenopathy. It is also known as necrotizing lymphadenitis. This is more common in adults but can occur in children with an average age of second to fourth decade of life. In adults, females are commonly affected and in paediatric age groups males are more commonly affected.<sup>1,2</sup> The etiology of Kikuchi disease is unknown but it can be triggered by some infections mostly viral and also by autoimmune diseases. Studies suggest that there is association with COVID-19 and Kikuchi disease.<sup>3,4</sup> Kikuchi disease is seen in patients with active COVID -19 and also may develop post covid. The Kikuchi disease is also seen in patients who have been administered with COVID -19 vaccination.<sup>5,6</sup> The usual symptoms of patients with Kikuchi disease are fever

with cervical lymphadenopathy. Very few cases present with generalized lymphadenopathy. We describe a young male presenting with multiple enlarged cervical lymph nodes. Fine needle aspiration needle showed a history of granulomatous lymphadenitis and Kikuchi-Fujimoto disease was diagnosed in histopathological examination and confirmed by immunohistochemistry.


## 2. Case History

A 19 year old male, came to the hospital with a history of fever and enlarged cervical lymph node. Ultrasound neck revealed bilateral cervical nodes with few matted, necrotic nodes. Fine needle aspiration of the lymph nodes revealed a cellular smear consisting of chronic inflammatory infiltrate of histiocytes, lymphocytes admixed with ill-defined granulomas consisting of epithelioid cells and lymphocytes in an extensive necrotic background. FNAC findings were suggestive of granulomatous lymphadenitis


\* Corresponding author.

E-mail address: [dr.manjani@bmch.ac.in](mailto:dr.manjani@bmch.ac.in) (S. Manjani).

and the smear was negative for acid fast stain. Other investigations such as sputum and gene-xpert studies were done to rule out tuberculosis. One lymph node was surgically removed and it was sent for histopathological examination. On gross examination, the node was 3 x 2 cm in size and externally capsulated, cut surface showed grey white areas. Microscopic sections revealed lymph node with fairly preserved architecture and reactive follicles composed of small lymphocytes admixed with immunoblasts in clusters and singly scattered. Focal areas of necrosis was seen along with histiocytes and nuclear debris. No definitive granuloma was made out and the sections were negative for Acid fast stain. The diagnosis of Kikuchi-Fujimoto disease was suggested and was confirmed by immunohistochemistry. Post-surgery the patient was medically managed and his condition improved well on regular follow up.



**Figure 1:** A: 40x Photomicrograph showing crescentic histiocytic cells in a granular background; B: 40x Photomicrograph showing necrotic debris; C: 4x-photomicrograph showing effaced node with alternating pale and dark areas.



**Figure 2:** A: 10x- Photomicrograph showing karyorrhectic debris with central necrosis; B: 40x- Photomicrograph showing phagocytic cell with eosinophilic cytoplasm and crescent shaped nuclei

### 3. Discussion

Kikuchi disease is one of the rare diseases which occurs most commonly in the South East Asian population. Among the affected patients the disease is common in young adults with a mean age of 20-30 years of age. Kikuchi disease is more common in females than in males with a female to male ratio of 2-4:1.<sup>7</sup>

Kikuchi disease occurs due to an unknown etiology but few studies show that Kikuchi disease can be triggered by viral infections or can be due to an autoimmune etiology. Few case reports suggest that Kikuchi disease can be triggered due to COVID -19 infection<sup>3,4</sup> and COVID -19 vaccinations.<sup>5,6</sup> The usual presentation of patients with Kikuchi disease will be acute in onset with complaints of fever, cervical lymphadenopathy which is usually unilateral.<sup>8,9</sup> Lymphadenopathy is observed in almost all the cases with Kikuchi disease and majority of the cases present with tender cervical lymphadenopathy and few cases may also have generalized lymphadenopathy.<sup>10</sup>

The diagnosis of Kikuchi disease is based on the histopathological findings followed by fine needle aspiration cytology. Possibility of tuberculosis and lymphoma must be considered and ruled out. Histological findings include preserved nodal architecture with areas of coagulative necrosis and absence of neutrophils.<sup>11</sup> Immunohistochemistry must be done to confirm the diagnosis of Kikuchi disease. The positive expression of immunohistochemical markers such as CD 68 and CD 4 suggests Kikuchi disease.<sup>12</sup>

The treatment of Kikuchi disease is supportive management along with steroid therapy. Though the disease resolves on its own, regular follow up must be done to know the condition of the patient and for disease recurrence.<sup>13</sup>

### 4. Conclusion

Kikuchi disease is rare self-limiting disease, though it is more common in eastern and western part of world. This disease should be always considered as differential diagnosis clinically and histomorphologically while reporting lymphoma in post pandemic era to avoid expensive molecular investigations

### 5. Source of Funding

None.

### 6. Conflict of Interest

None.

### References

1. Joean O, Thiele T, Raap M, Schmidt RE, Stoll M. Take a second look: it's Kikuchi's disease! A case report and review of literature. *Clinics*

- and Practice. 2018;8(4):111–3. doi:10.4081/cp.2018.1095.
2. Perry AM, Choi SM. Kikuchi-Fujimoto Disease: A Review. *Arch Pathol Lab Med*. 2018;142(11):1341–6.
  3. Jaseb K, Fard NN, Rezaei N, Sadeghian S, Sadeghian S. COVID-19 in a case with Kikuchi-Fujimoto disease. *Clin Case Rep*. 2021;9(3):1279–82.
  4. Ghadeer HA, AlKadhem SM, AlMajed MS, AlAmer HM, AlHabeeb JA, Alomran SH, et al. Kikuchi-Fujimoto disease following COVID-19. *Cureus*. 2009;14(1):e21049. doi:10.7759/cureus.21049.
  5. Soub H, Ibrahim W, Maslamani MA, Maslamani M, Ali GA, Ummar W, et al. Kikuchi-Fujimoto disease following SARS CoV2 vaccination: Case report. *IDCases*. 2021;25:01253. doi:10.1016/j.idcr.2021.e0125.
  6. Tan HM, Hue SS, Wee A, See KC. Kikuchi-Fujimoto Disease Post COVID-19 Vaccination: Case Report and Review of Literature. *Vaccines*. 2021;9(11):1251. doi:10.3390/vaccines9111251.
  7. Dumas G, Fain O. Kikuchi-Fujimoto Disease. In: A Clinician's Pearls & Myths in Rheumatology. Cham: Springer International Publishing; 2023. p. 743–7.
  8. Harrison J, Sukumaran S, Vijayan V. Systemic Lupus Erythematosus Lymphadenopathy Presenting as Kikuchi-Fujimoto Disease in an Adolescent. *Cureus*. 2023;15(2):e35304. doi:10.7759/cureus.35304.
  9. Baek JY, Kang JM, Lee JY, Lim SM, Ahn JG. Comparison of Clinical Characteristics and Risk Factors for Recurrence of Kikuchi-Fujimoto Disease Between Children and Adult. *J Inflamm Res*. 2022;15:5505–14.
  10. Kellner D, Temelkovska T, Greeley A, Saito A. Kikuchi-Fujimoto Disease: A Case Report of Fever and Lymphadenopathy in a Young White Man. *Fed Pract*. 2023;40(1):22–7.
  11. Song Y, Liu S, Song L, Chen H, Bai M, Yan J, et al. Case report: histiocytic necrotizing lymphadenitis (Kikuchi-Fujimoto Disease) concurrent with aseptic meningitis. *Front Neurol*. 2021;12:565387. doi:10.3389/fneur.2021.565387.
  12. Jagadeesan S, Rani S, Porwal YC, Patel P. Lupus and Kikuchi-Fujimoto Disease: A Combination for Catastrophe. *Cureus*. 2022;14(7):e27423. doi:10.7759/cureus.27423.
  13. Semavi DG, Turgay UL. Highlighting steroid indications for Kikuchi Fujimoto disease: A case report. *Cukurova Med J*. 2022;47(4):1753–5.

## Author biography

**Jackson Bose**, Assistant Professor

**Jamunarani Srirangaramasamy**, Professor and HOD

**Khowsalya Subrajaa Karupasamy**, Assistant Professor

**Sekar Manjani**, Associate Professor

**Cite this article:** Bose J, Srirangaramasamy J, Karupasamy KS, Manjani S. Kikuchi-Fujimoto disease in a young male post-COVID -19 vaccination. *IP Arch Cytol Histopathology Res* 2023;8(4):268-270.