

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Archives of Cytology and Histopathology Research

Journal homepage: <https://www.achr.co.in/>

Case Report

A rare case report of hurthle cell adenoma mimicking malignancy

Shashikala Gurappa Hiremani^{1*}

¹Dept. of Pathology, District Hospital, Vijayapura, Karnataka, India



ARTICLE INFO

Article history:

Received 17-07-2024

Accepted 22-08-2024

Available online 08-10-2024

Keywords:

Thyroid lesions

Hurthle cell tumors

FNAC

IHC

ABSTRACT

Introduction: Thyroid lesions are the most common endocrine diseases, occurring in usually all age groups, though accurate diagnosis still remains a challenge, as due to oncocytic potentials of Thyroid cells leads to like indeterminate or suspicious diagnosis. Long standing benign Thyroid lesion shows atypical presentation suspicious to malignancy. Now a days with the help of WHO classification and available advanced diagnostic measures helping to confirm the diagnosis, avoiding overdiagnosis and wrong treatment. Hence this study review the importance of Histopathological study with coordination with immunohistochemistry (IHC) study, where suspicious thyroid malignant lesion finally diagnosed as benign Hurthle cell/ Oncocytic adenoma in Nodular Goitre.

Case Report: 72 year old female presented huge neck swelling since one year, was reviewed in our departments of Pathology and ENT. Fine needle aspiration cytology (FNAC) features are suspicious of Malignancy, further Histopathological features were studied with co ordance of IHC study.

Results: FNAC features, which were suspicious of Malignancy were further studied with Histopathological and IHC studies diagnosed as Hurthle cell/ Oncocytic adenoma in Nodular Goitre.

Conclusion: Histopathology remains the cornerstone for the diagnosis of thyroid oncocytic lesions and also in distinguishing between adenoma and carcinoma.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Thyroid lesions are the most common endocrine diseases, with nodular architecture and follicular pattern of growth often pose difficulties in accurate diagnosis during the assessment of cytologic and histologic specimens.¹ Majority of Thyroid tumours originates from follicular epithelial cells.

Hurthle cell adenomas are rare, benign tumours of the thyroid, accounting for less than 5%. Size more than 4cms rare.²

Hurthle cell adenomas are noninvasive, encapsulated tumors which are surrounded by a thin capsule and

characterized by cells with abundant granular, eosinophilic cytoplasm with prominent central nucleoli.

These tumors are remain interesting diagnostic phenomena with acknowledged preoperative and intraoperative adversities in discriminating adenomas from carcinomas.

Hurthle cell tumors often challenging to diagnose due to the ubiquity of Hurthle cells ranging from non-neoplastic conditions to aggressive malignancies. Galectin-3 is a human lectin linked to malignant transformation in different organs including thyroid gland.³

2. Aim of Study

To project the importance of histopathological examination in distinguishing between adenoma and carcinoma.

* Corresponding author.

E-mail address: shashikalagurappahiremani@gmail.com (S. G. Hiremani).

3. Case Report

3.1. History

A 72-year-old woman presented with midline diffuse Thyroid swelling in neck since one year, as shown in figure 1. She had no symptoms of weight loss, pain and pressure, hoarseness or fatigue. Her neck swelling was insidious in onset, gradually progressive, initially a size of 2x1 cms and progressed to size of 10x8cms. Physical examination of neck showed a solitary swelling of size 10x8cms present in front of neck more towards right side and moved with deglutition.



Figure 1: Diffuse Thyroid swelling in neck.

3.2. Physical examination of neck

A solitary swelling of size 12x8 cms present in front of neck more towards right side and moved with deglutition.

3.3. Investigations

Routine Laboratory indices were within normal limits.

3.3.1. Ultrasound of neck

Showed thick walled heteroeic lesion occupying whole of right lobe with a TI-RADS score - Four suggesting Suspicious for malignancy.




Figure 2: FNAC– showing large polygonal cells with pleomorphic features.

3.3.2. FNAC of thyroid swelling

Showed high cellularity comprised of large polygonal cells with nuclear pleomorphism, prominent nucleoli and was reported as SUSPICIOUS FOR MALIGNANCY (The Bethesda System of Reporting Thyroid Cytopathology-V) as in Figure 2.

3.3.3. Histopathology study

Total thyroidectomy was done, specimen was sent for histopathological study.

Grossly the mass is as shown in Figures 3 and 4.

Microscopy revealed a well encapsulated tumor showing oncocytic cells with abundant granular and eosinophilic cytoplasm arranged in micro and macrofollicular pattern, at places papillary pattern. Peripheral thyroid tissue showed features of Nodular Goitre. No evidence of vascular/capsular invasion was seen in around 60 sections after extensive grossing.

A diagnosis of Hurthle cell/ Oncocytic adenoma in Nodular Goitre was rendered on histopathology as shown in Figures 5 and 6.




Figure 3: Total thyroidectomy specimen. Right lobe, measuring 8x7x4 cm. dark brown smooth glistening. Left lobe 4x2x1cm.

3.3.4. Immunohistochemistry (IHC) study

The tumor was negative for IHC marker Galectin Figure 7, ruling out Papillary carcinoma of thyroid. However IHC for Ki 67 showed a moderate mitotic activity of 24-26%.

Final diagnosis: Hurthle cell/ Oncocytic adenoma in Nodular Goitre.




Figure 4: Cut section—shows multiple friable brown tissue bits and drained 20-30ml of colloid.




Figure 5: 20x H and E stain –Showing tumor cells arranged in micro and macrofollicular pattern and focal papillary pattern.




Figure 6: 40x H and E stain - Showing oncocyctic cells with abundant granular eosinophilic cytoplasm.




Figure 7: IHC for Galectin -3 negative.

4. Discussion

When compared with classic PTC, DSV has unique clinical features, including a higher prevalence of underlying Hashimoto's thyroiditis, higher female-to-male ratio, and younger age⁴, but in our study, patient is elderly female.

Our case presented with only palpable huge neck mass. As size more than 4cms rare,² as similarly in our case, the swelling was 10x8cms, hence our rare entity . Usually thyroid nodules present along with palpable neck mass, anterior neck pain, globus sensation, dysphagia, dyspnea, and dysphonia/ hoarseness.⁵

Till date as similar to our case, poorly differentiated thyroid carcinoma remains a challenge due to its aggressive disease course.⁶

Detection rate of thyroid nodules varies widely in population from 4% to 67%, by physical examination around 4-7%, with a higher detection rate of 30-67% by ultrasound.⁷⁻⁹ but in our case patient noticed huge increased in swelling size and came to hospital.

Once the results of a thyroid FNAC are available, the risk of thyroid cancer can be estimated and can guide the next steps in management in patients with benign or malignant results,^{10,11} similarly as in our study FNAC report was suspicious for malignancy (the Bethesda system of reporting thyroid cytopathology-v) guided for further Histopathology and IHC studies.

Hürthle cell adenoma is a rare variant of benign lesion of thyroid gland, however aggressive nature and potential controversies about its malignant nature still remains.

Hürthle cell tumour though rare, yet they are often challenging to diagnose accurately with FNAC, with significant interobserver variability, largely due to the ubiquity of Hürthle cells ranging from nonneoplastic conditions to aggressive malignancies,¹² as in our study FNAC reported as suspicious for malignancy.

Hürthle cell tumors have been associated with various of benign thyroid disorders, including Hashimoto's thyroiditis, hyperthyroidism, nodular goitre, and thyroid neoplasms,

where they are thought to be oncocytic metaplasia,¹³ as in our study the FNAC report suspicious of malignancy, might be due to potentials of oncocytic metaplasia.

Pre-operative diagnosis is difficult, fine needle aspiration cytology may show Hürthle cell lesions, but final diagnosis is confirmed by histopathology for capsular and vascular invasion,¹⁴ as similarly in our case, FNAC features of suspicious for malignancy and final diagnosis confirmed by histopathology and IHC negative for Galectin-3 study, as retaining importance of histopathology and IHC study as goldstandard investigations for correct diagnosis.

As of the most recent WHO classification, thyroid tumours are a form of tumour that arises from thyroid follicles,¹⁵ as similar in our case, tumour arised from thyroid origin.

As similar studies,^{16,17} ultrasound preoperative diagnosis is difficult. Hürthle cell cancer (HCC) appears as hypochoic to hyperechoic lesions on ultrasonography, but in our study, thyroid swelling appeared as heteroeoic lesion on ultrasonography.

Similar to in our study, majority of Thyroid neoplams the intraoperative frozen section and FNAC cannot reliably differentiate cancer from adenoma.¹⁸

Hurthle cell tumors are distinct, not only because of the enduring debate about the true nature, but also because HCC are particularly aggressive, thereby emphasizing the need to differentiate Hurthle cell adenomas from carcinomas.

Hürthle cell carcinoma has a more aggressive clinical course than any other differentiated thyroid carcinoma, with particular drawbacks such as a greater rate of regional lymph node metastases, recurrences, and cause-specific fatality,¹⁹ hence FNAC report was suspicious, hence further proceeded to Histopathology and IHC study.

The differential diagnosis include Hashimoto's thyroiditis, Nodular goitre, Medullary carcinoma with oncocytic variant and Papillary thyroid carcinoma with oncocytic variant, warthin-like variant and tall cell variant.

The ongoing elucidation of a malignant HCC depends on capsular and/or vascular invasion, and the presence of metastasis. This cannot be determined by cytologic evaluation and relies on histological examination of a resected specimen.

In our case, although being a benign tumor on histopathology and IHC, showed features of malignancy on imaging, cytology and on frozen section.

In addition, Gholami et al.²⁰ pointed out the FNAC was performed for all the nodules and 39 (70.9%) were benign and 16 (29.1%) were malignant or suspicious for malignancy, were referred for surgery, similarly in our case, for suspicious for malignancy as per FNAC report, total Thyroidectomy was done for further evaluation and diagnosis.

5. Conclusion

Immunohistochemistry is the most commonly used method to complement morphologic assessment.

Histopathology remains the cornerstone for the diagnosis of thyroid oncocytic lesions and also in distinguishing between adenoma and carcinoma.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

1. Fischer SE. Application of immunohistochemistry to thyroid neoplasms. *Arch Pathol Lab Med.* 2008;132(3):359–72.
2. Singh RK, Goyal A, Elhence P, Kumar A. Bilateral Hurthle cell adenoma of thyroid: a rare case report. *Ijorlcom cited.* 2017;3(2):462–4.
3. Aiad HA, Kandil MA, Asaad NY, El-Kased AM, El-Goday SF. Galectin-3 immunostaining in cytological and histopathological diagnosis of thyroid lesions. *J Egypt Natl Canc Inst.* 2008;20(1):36–46.
4. Chow SM, Chan JKC, Law SCK, Tang DLC, Ho CM, Cheung WY, et al. Diffuse sclerosing variant of papillary thyroid carcinoma-clinical features and outcome. *Eur J Surg Oncol.* 2003;29(5):446–9.
5. Sørensen JE, Hegedüs L, Kruse-Andersen S, Kruse-Andersen S, Bonnema SJ. The impact of goitre and its treatment on the trachea, airflow, oesophagus and swallowing function. A systematic review. *Best Pract Res Clin Endocrinol Metab.* 2014;28(4):481–94.
6. Agarwal A, Nelson G, Niraj K, Narendra K, Prabhaker M, Sushil G, et al. Poorly differentiated thyroid cancer: Clinical, pathological, mutational, and outcome analysis. *Indian J Pathol Microbiol.* 2024;doi:10.4103/ijpm.ijpm_885_23.
7. Guth S, Theune U, Aberle J, Galach A, Bamberger CM. Very high prevalence of thyroid nodules detected by high frequency (13 MHz) ultrasound examination. *Eur J Clin Invest.* 2009;39(8):699–706.
8. Reiners C, Wegscheider K, Schicha H, Theissen P, Vaupel R, Wrbitzky R, et al. Prevalence of thyroid disorders in the working population of Germany: ultrasonography screening in 96,278 unselected employees. *Thyroid.* 2004;14(11):926–32.
9. Tufano RP, Noureldine SI, Angelos P. Incidental thyroid nodules and thyroid cancer: considerations before determining management. *JAMA Otolaryngol Head Neck Surg.* 2015;141(6):566–72.
10. Haugen BR, Alexander EK, Bible KC, Doherty GM, Mandel SJ, Nikiforov YE, et al. American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid.* 2015;26(1):1–133.
11. Ospina NS, Brito JP, Maraka S, Ycaza AEE, Rodriguez-Gutierrez R, Gionfriddo MR, et al. Diagnostic accuracy of ultrasound-guided fine needle aspiration biopsy for thyroid malignancy: systematic review and meta-analysis. *Endocrine.* 2016;53(3):651–61.
12. Wong KS, Angell TE, Barletta JA, Krane JF. Hürthle cell lesions of the thyroid: Progress made and challenges remaining. *Cancer Cytopathol.* 2021;129(5):347–62.
13. Singh RK, Goyal A, Elhence P, Kumar A. Bilateral Hurthle cell adenoma of thyroid: a rare case report. *Int J Otorhinolaryngol Head Neck Surg.* 2017;3(2):462–4.
14. Tambat RM. Hürthle Cell Adenoma in a Hashimoto's Thyroiditis Patient: A Rare Case Report. *Clin Case Rep.* 2022;4(3):194. doi:10.46527/2582-5038.194.
15. Lloyd RV, Osamura RY, Klöppel G, Rosai J. WHO Classification of Tumours of Endocrine Organs. 4th ed. France: IARC Press; 2017.

16. Santana NO, Freitas RMC, Marcos VN, Chammas MC, Camargo RYA, Schmerling CK, et al. Diagnostic performance of thyroid ultrasound in Hürthle cell carcinomas. *Arch Endocrinol Metab.* 2019;63(3):300–5.
17. Maizlin ZV, Wiseman SM, Vora P, Kirby JM, Mason AC, Filipenko D, et al. Hürthle cell neoplasms of the thyroid: sonographic appearance and histologic characteristics. *J Ultrasound Med.* 2008;27(5):751–7.
18. Chen H, Nicol TL, Udelsman R. Follicular lesions of the thyroid. Does frozen section evaluation alter operative management? *Ann Surg.* 1995;222(1):101–6.
19. Haugen BR, Sawka AM, Alexander EK, Bible KC, Caturegli P, Doherty GM. American Thyroid Association Guidelines on the Management of Thyroid Nodules and Differentiated Thyroid Cancer Task Force Review and Recommendation on the Proposed Renaming of Encapsulated Follicular Variant Papillary Thyroid Carcinoma Without Invasion to Noninvasive Follicular Thyroid Neoplasm with Papillary-Like Nuclear Features. *Thyroid.* 2017;27(4):481–3.
20. Gholami M, Aghaghazvini L, Gholampour M, Kargar J. Evaluation of the diagnostic value of Ultrasound Elastography in TIRADS 4 categories of Thyroid nodules". *J Adv Pharm Edu Res.* 2021;11(S1):26–32.

Author biography

Shashikala Gurappa Hiremani, Senior Pathologist and HOD

Cite this article: Hiremani SG. A rare case report of hurthle cell adenoma mimicking malignancy. *IP Arch Cytol Histopathology Res* 2024;9(3):158-162.